

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.
- 2 Simplify problem: drop the absolute values and maximize $(x_i - x_j) - c(i - j) = (x_i - c \cdot i) + (-x_j + c \cdot j)$.

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.
- 2 Simplify problem: drop the absolute values and maximize $(x_i - x_j) - c(i - j) = (x_i - c \cdot i) + (-x_j + c \cdot j)$.
 - Keep track of largest value y of $-x_j + c \cdot j$ seen so far.

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.
- 2 Simplify problem: drop the absolute values and maximize $(x_i - x_j) - c(i - j) = (x_i - c \cdot i) + (-x_j + c \cdot j)$.
 - Keep track of largest value y of $-x_j + c \cdot j$ seen so far.
 - Answer to simplified problem is $x_i - c \cdot i + y$.

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.
- 2 Simplify problem: drop the absolute values and maximize $(x_i - x_j) - c(i - j) = (x_i - c \cdot i) + (-x_j + c \cdot j)$.
 - Keep track of largest value y of $-x_j + c \cdot j$ seen so far.
 - Answer to simplified problem is $x_i - c \cdot i + y$.
- 3 To solve original problem, separately solve variant for $(x_j - x_i) - c(j - i)$ the same way and take largest of the two.

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.
- 2 Simplify problem: drop the absolute values and maximize $(x_i - x_j) - c(i - j) = (x_i - c \cdot i) + (-x_j + c \cdot j)$.
 - Keep track of largest value y of $-x_j + c \cdot j$ seen so far.
 - Answer to simplified problem is $x_i - c \cdot i + y$.
- 3 To solve original problem, separately solve variant for $(x_j - x_i) - c(j - i)$ the same way and take largest of the two.
- 4 Time complexity $O(n)$.

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.
- 2 Simplify problem: drop the absolute values and maximize $(x_i - x_j) - c(i - j) = (x_i - c \cdot i) + (-x_j + c \cdot j)$.
 - Keep track of largest value y of $-x_j + c \cdot j$ seen so far.
 - Answer to simplified problem is $x_i - c \cdot i + y$.
- 3 To solve original problem, separately solve variant for $(x_j - x_i) - c(i - j)$ the same way and take largest of the two.
- 4 Time complexity $O(n)$.
- 5 More complicated solutions using balanced search trees or range trees also possible.

A — Antenna Analysis

Problem

Given integers x_1, \dots, x_n , find, for each i , the maximum of $|x_i - x_j| - c|i - j|$ over $j \leq i$.

Solution

- 1 Key observation: dropping the absolute values does not increase the answer.
- 2 Simplify problem: drop the absolute values and maximize $(x_i - x_j) - c(i - j) = (x_i - c \cdot i) + (-x_j + c \cdot j)$.
 - Keep track of largest value y of $-x_j + c \cdot j$ seen so far.
 - Answer to simplified problem is $x_i - c \cdot i + y$.
- 3 To solve original problem, separately solve variant for $(x_j - x_i) - c(i - j)$ the same way and take largest of the two.
- 4 Time complexity $O(n)$.
- 5 More complicated solutions using balanced search trees or range trees also possible.

Statistics at 4-hour mark: 626 submissions, 88 accepted, first after 00:04