

---

## Problem A. Easy Problem

Input file:            **standard input**  
Output file:         **standard output**  
Time limit:          2 seconds  
Memory limit:       512 megabytes

A sequence  $(a_1, a_2, \dots, a_n)$  is  $(n, m, d)$ -good if  $1 \leq a_i \leq m$  ( $1 \leq i \leq n$ ) and  $\gcd(a_1, a_2, \dots, a_n) = d$ .

Given four integers  $n$ ,  $m$ ,  $d$  and  $k$ , you are asked to calculate the sum of  $f(q, k)$  for each  $(n, m, d)$ -good sequence  $q$ , where  $f((a_1, a_2, \dots, a_n), k) = (a_1 a_2 \dots a_n)^k$  for the sequence  $q = (a_1, a_2, \dots, a_n)$ .

Since the answer could be very large, you only need to output the answer modulo 59964251.

### Input

The first line is an integer  $T$  ( $1 \leq T \leq 20$ ), which is the number of test cases.

For each test case, the first line contains four integers  $n$  ( $1 \leq n \leq 10^{100000}$ ),  $m$  ( $1 \leq m \leq 100000$ ),  $d$  ( $1 \leq d \leq 100000$ ),  $k$  ( $1 \leq k \leq 10^9$ ), which are described in the problem description.

### Output

For each test case, output a line containing a single integer.

### Example

standard input	standard output
1 3 3 3 1	27