

Operator Precedence

Input file: *standard input*
Output file: *standard output*
Time limit: 1 second
Memory limit: 1024 mebibytes

Randias is facing his primary school homework:

Find a nonzero integer sequence a of length $2n$ satisfying

$$(a_1 \times a_2) + (a_3 \times a_4) + \dots + (a_{2n-1} \times a_{2n}) \\ = a_1 \times (a_2 + a_3) \times (a_4 + a_5) \times \dots \times (a_{2n-2} + a_{2n-1}) \times a_{2n} \neq 0.$$

In shorter form, $\sum_{i=1}^n a_{2i-1} a_{2i} = a_1 a_{2n} \prod_{i=2}^n (a_{2i-2} + a_{2i-1}) \neq 0$.

Of course, Randias knows how to solve it. But he wants to give you a test. Can you solve the question above?

Input

Each test contains multiple test cases. The first line contains a single integer t ($1 \leq t \leq 10^5$) denoting the number of test cases.

For each test case, the only line contains a single integer n ($2 \leq n \leq 10^5$).

It is guaranteed that the sum of n over all test cases does not exceed $2 \cdot 10^5$.

Output

For each test case, output one line with $2n$ integers: a_1, a_2, \dots, a_{2n} ($1 \leq |a_i| \leq 10^{10}$).

It can be shown that the answer always exists.

If there are several possible answers, output any one of them.

Example

<i>standard input</i>	<i>standard output</i>
3	1 -3 -3 1
2	1 -10 6 6 -10 1
3	1 -15 10 -1 -1 10 -15 1
4	